



ATLANTIK™

Hand Harvested, Concentrated Extracts Of *Ascophyllum nodosum*



9 341200 000607

20 L

ATLANTIK DIRECTIONS FOR USE

SITUATION	CROPS	RATE	COMMENTS
<i>Broadacre</i>	Wheat , Barley, Sorghum, Oats, Rice, Cotton, Maize	500mL/ha	Apply at 4 to 6 leaf stage
		1L/ha	Apply at boot stage/pre flower or during times of stress
<i>Brassicas</i>	Canola, Broccoli, Cauliflower, Cabbage	750mL/ha	Apply at 3 to 4 true leaf stage
		1L/ha	Apply at flowering or head formation and during times of stress
<i>Bulb crops</i>	Garlic, Shallots, Onions	1L/ha	Apply at bulb initiation followed by applications every 10 to 14 days if required or during times of stress
<i>Legumes</i>	Beans, Peas, Chickpea, Peanuts	500mL/ha	Apply at 3 to 5 trifoliolate or 6 weeks after emergence
		1L/ha	Apply pre flower and at pegging or during times of stress
<i>Cucurbits</i>	Squash, Pumpkin, Rockmelon, Honeydew, Watermelon, Zucchini, Cucumber	500mL/ha	Apply at 3 to 4 true leaf stage
		1L/ha	Apply at commencement of runner stage and at pre flower or during times of stress
<i>Leafy vegetables</i>	Celery, Lettuce, Wombok, Herbs	750mL/ha	Apply at plant establishment
		1L/ha	Apply every 7 to 10 days as required or during times of stress
<i>Vegetables</i>	Capsicum, Peppers, Eggplant, Tomato	500mL/ha	Apply at plant establishment
		1L/ha	Apply every 7 to 10 days as required or during times of stress
<i>Berries</i>	Strawberry, Blueberry, Raspberry	1L/ha	Apply at bud break
		1.5L/ha	Apply at 14 day intervals as required or during times of stress
<i>Tree Crops</i>	Fruit	1L/ha	Apply at bud break / pre flower then apply at petal fall
		2L/ha	Apply during fruit fill/development or during times of stress
	Nuts	1L/ha	Apply at pre flowering
		2L/ha	Apply during nut fill/development as required or during times of stress
<i>Vines</i>	Grapes, Passionfruit, Kiwifruit	1L/ha	Apply at bud break then at pre flower as required
		2L/ha	Apply during fruit fill/development or during times of stress at a minimum of 14 day intervals
<i>Sugarcane</i>	Plant cane	2L/ha	Apply in furrow over cane sett prior to cover up
			Apply during early growth ensuring adequate coverage of foliage
	Ratoon cane		Apply when sufficient foliage available ensuring adequate coverage
<i>Root/Tuber</i>	Carrots, Potatoes, Beetroot	750mL/ha	Apply at 4 to 6 leaf stage
		1L/ha	Apply when stolons are developing then repeat at tuber bulking if required or during times of stress
<i>Transplants</i>	Dipping	1 : 250	Dip transplants /cane sett/propagules in solution until plants are thoroughly wet
	Seedling Tray Drench	6mL/L	Drench seedling trays with solution using watering can then allow to dry before attempting to remove seedlings from tray

APPLICATION INSTRUCTIONS: Shake well before using.

Apply with sufficient water to achieve thorough cover of foliage. Apply up to 50L/ha by aircraft and up to 2000 L/ha for ground applied equipment. Do not exceed a dilution of 1L of Atlantik with 50 L of water.

GENERAL COMMENTS

DO NOT apply Atlantik with organosilicone type surfactants or other cuticle disruptive adjuvants.

DO NOT mix Atlantik with Calcium products other than Calcium EDTA.

DO NOT mix Atlantik with highly acidic solutions or strong oxidizing agents as product performance may be altered. To improve rainfastness, apply with Hygro-Stic[®], Flexstic[®] or Flexlend[®] as per directions for each product label.

MIXING INSTRUCTIONS:

Shake well before use. Add Atlantik to a half filled spray tank of water, then continue adding water while agitating. Continue agitation until the product is completely mixed.

COMPATIBILITY

Compatible with most commonly used fertilizers and agrochemicals. It is recommended not to mix Atlantik with calcium products other than Calcium EDTA.

It is not recommended to mix Atlantik with strong acidic nor strong oxidising agents. You must perform a jar test before mixing commercial quantities.

STORAGE: Store in the closed original container, in a cool, well-ventilated area away from children, pets, animals, food and feedstuffs. Do not store for prolonged periods in direct sunlight.

WARNING: Keep out of reach of children. If product comes into contact with skin or eyes, wash immediately with plenty of water. Store in a fresh, dry place and protect from excessive heat or cold. Store the product in its original container. Use the product according to the instructions on the label. If in doubt, please contact your local agronomist. If local experience is unavailable, it is recommended a small-scale test application be conducted before widespread use.

USE ONLY IN CASE OF PROVEN NECESSITY.

DO NOT EXCEED THE MAXIMUM DOSAGES.

CONDITION OF SALE: AGSPEC warrants that the product within the unopened pack conforms to the chemical description on this label. As the occurrence of unusual environmental conditions, and the use of this product, is all beyond the control of AGSPEC, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of AGSPEC exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of the manufacturer, available on request. Use of the product implies that the purchaser accepts these conditions of sale.



Packed and Distributed in Australia by
Agspec Australia Pty Ltd.
ABN: 401 095 739 53
PO. Box 1006, Mt. Gambier, S.A., 5290
Mobile - 0427 490 551

Atlantik™ is a trademark of Agspec Australia Pty. Ltd.

Hygro-Stic[®], Flexstic[®] or Flexlend[®] are registered trademarks of Agspec Australia Pty. Ltd.



Product of Ireland.
Extracted using patented
extraction optimisation process (EOP)

